

Total No. of Printed Pages—7

**1 SEM TDC MTMH (CBCS) C 2**

**2 0 2 2**

( Nov/Dec )

**MATHEMATICS**

( Core )

Paper : C-2

( Algebra )

Full Marks : 80

Pass Marks : 32

Time : 3 hours

*The figures in the margin indicate full marks  
for the questions*

1. (a) State the modulus of the complex number  $(1 + \cos\theta + i\sin\theta)^5$ . 1

(b) If  $\cos\alpha + \cos\beta + \cos\gamma = 0$   
 $\qquad\qquad\qquad = \sin\alpha + \sin\beta + \sin\gamma$

then show that

$$\sin 3\alpha + \sin 3\beta + \sin 3\gamma = 3 \sin(\alpha + \beta + \gamma) \quad 2$$

(c) Show that

$$(1+i)^n + (1-i)^n = 2^{\frac{n}{2}+1} \cos \frac{n\pi}{4} \quad 3$$

Or

If  $\text{cis}\theta = \cos\theta + i\sin\theta$  and  $x = \text{cis}\alpha$ ;  
 $y = \text{cis}\beta$ ;  $z = \text{cis}\gamma$  with  $x + y + z = 0$ , show  
 that  $x^{-1} + y^{-1} + z^{-1} = 0$ .

(d) Show that the product of  $n$ -numbers of  
 $n$ th root of unity is  $(-1)^{n-1}$  and their sum  
 is zero. 4

2. (a) Explain why the set of integers with the  
 relation 'less than or equal to' ( $\leq$ ) is not  
 an equivalence relation. 1

(b) Give an example of a bijective map. 1

(c) Given  $f(x) = |x|$ , show that  
 $(f \circ f)(x) = f(x)$ . 2

(d) If  $\text{g.c.d}(a, b) = d$ , show that

$$\text{g.c.d.} \left( \frac{a}{d}, \frac{b}{d} \right) = 1 \quad 2$$

(e) Show that the relation of equality on the set of integers is an equivalence relation. 3

(f) Use mathematical induction to show that (any one)—

(i) 2 is a factor of  $5^n - 3^n \forall n \in \mathbb{N}$ ;

(ii)  $1^3 + 2^3 + \dots + n^3 = \left[ \frac{n}{2}(n+1) \right]^2$ . 3

(g) Show that if  $f: X \rightarrow Y$  is a bijection, then  $\exists$  a map  $g: Y \rightarrow X$  such that  $g \circ f$  and  $f \circ g$  are identity maps. 3

(h) Let  $k > 0$  be an integer and  $j$  be any integer. Then show that  $\exists$  unique integers  $q$  and  $r$  such that  $j = kq + r$  where  $0 \leq r < k$ . 5

(i) Show that if  $a$  is an odd integer, then  $a^{2^n} \equiv 1 \pmod{2^{n+2}}$  for any  $n \geq 1$ . 5

3. (a) State whether true or false : 1

Each matrix is row equivalent to one and only one reduced Echelon matrix.

- (b) Fill in the blank : 1

The equation  $x = \alpha u + \beta v$  where  $\alpha$  and  $\beta$  are fixed scalars and neither  $u$  nor  $v$  is a multiple of the other, geometrically represents \_\_\_\_\_ through the origin.

- (c) Solve

$$x_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

and state the nature of the two non-zero vectors. 1+1=2

- (d) State whether the following vectors are linearly dependent or independent by inspection justifying the reason thereof :

1+1=2

$$\begin{bmatrix} 3 \\ 5 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 6 \\ 5 \\ 4 \end{bmatrix}$$

- (e) Show that  $\forall u, v, w \in \mathbb{R}^n$ ,  
 $(u+v)+w = u+(v+w)$ . 2

- (f) Reduce the matrix

$$\begin{bmatrix} 1 & 3 & 1 \\ -4 & -9 & 2 \\ 0 & -3 & -6 \end{bmatrix}$$

to row reduced Echelon form using forward and backward phases of row operations. 4

- (g) Solve the following system by reducing the augmented matrix to row reduced Echelon form indicating the basic and free variables : 4

$$\begin{aligned} x_1 + 3x_2 - 5x_3 &= 4 \\ x_1 + 4x_2 - 8x_3 &= 7 \\ -3x_1 - 7x_2 + 9x_3 &= -6 \end{aligned}$$

- (h) For an  $m \times n$  matrix  $A$ , if  $u, v \in \mathbb{R}^n$ , and  $c$  is any scalar, show that—

(i)  $A(u + v) = Au + Av$ ;

(ii)  $A(cu) = c(Au)$ . 2+2=4

4. (a) For a linear transformation  $T$ , show that  $T(0) = 0$ . 1

- (b) For the linear transformation  $T: \mathbb{R}^5 \rightarrow \mathbb{R}^2$  given by  $T(x) = Ax$ , state the order of the matrix  $A$ . 1

- (c) For  $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , give the geometric description of the transformation  $x \mapsto Ax$ . 2
- (d) Show that the map  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = mx$  is a linear transformation. 2
- (e) Let  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  be linear. Show that  $\exists$  a unique matrix  $A$  such that  $T(x) = Ax$   $\forall x \in \mathbb{R}^n$ . 3
- (f) If  $A$  is an invertible  $n \times n$  matrix, then  $\forall b \in \mathbb{R}^n$ , show that the matrix equation  $Ax = b$  has the unique solution  $x = A^{-1}b$ . 3
- (g) Show that null  $A$  is a subspace of  $\mathbb{R}^n$ . 4
- (h) Find the eigenvalues of
- $$A = \begin{bmatrix} 4 & 0 & 1 \\ -2 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$
- 4

( 7 )

- (i) Find the bases for col A and null A stating their dimensions where

$$A = \begin{bmatrix} 1 & -3 & 2 & -4 \\ -3 & 9 & -1 & 5 \\ 2 & -6 & 4 & -3 \\ -4 & 12 & 2 & 7 \end{bmatrix}$$

5

\*\*\*