3 SEM TDC MTMH (CBCS) C 7

2021

(Held in January/February, 2022)

MATHEMATICS

(Core)

Paper: C-7

(PDE and Systems of ODE)

Full Marks: 60
Pass Marks: 24

Time: 3 hours

The figures in the margin indicate full marks for the questions

1. (a) Write the degree of the equation

$$x\left(\frac{\partial^2 z}{\partial x^2}\right) + \left(\frac{\partial z}{\partial y}\right)^2 = \frac{\partial z}{\partial x}$$

(b) Write Lagrange's subsidiary equation of

xzp + yzq = xy

(c) Write the general solution of pq = k.

(d) Solve: 5

$$(y-zx)p+(x+yz)q=x^2+y^2$$

Or

Find the integral surface of $x^2p+y^2q+z^2=0$, which passes through the hyperbola xy=x+y, z=1.

(e) Show that the equations xp - yq = x and $x^2p + q = xz$ are compatible.

(Turn Over)

1

1

1

5

22P/93

- **2.** (a) Write Charpit's auxiliary equations for $q = 3p^2$.
 - (b) Find complete integral of any one of the following:

(i)
$$q = (z + px)^2$$

(ii)
$$q + px = p^2$$

(iii)
$$z^2 = pqxy$$

(c) Find a complete integral of

$$p_1^3 + p_2^2 + p_3 = 1$$

Or

Solve the boundary value problem $\frac{\partial u}{\partial x} = 4 \frac{\partial u}{\partial y}$ with $u(0, y) = 8e^{-3y}$ by the method of separation of variables.

3. (a) Write the condition when the equation Rs + Ss + Tt + f(x, y, z, p, q) = 0

is hyperbolic.

(b) Determine the nature of the equation

$$\frac{\partial^2 z}{\partial x^2} = x^2 \frac{\partial^2 z}{\partial y^2}$$

(c) Show that u = f(x+y) + g(y-x) satisfies the equation

$$\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} = 0$$

where f and g are functions.

2

4

1

2

2

(d) Reduce the equation $\frac{\partial^2 z}{\partial x^2} + x^2 \frac{\partial^2 z}{\partial y^2} = 0$ to canonical form.

7

1

Or

Derive the one-dimensional heat equation.

- **4.** (a) Write the general form of two-dimensional heat equation.
 - (b) Write one assumption on vibrating string problem.
 - (c) Solve

$$\frac{\partial^2 u}{\partial x^2} = k^2 \left(\frac{\partial u}{\partial t} \right)$$

when u(0, t) = u(l, t) = 0, $u(x, 0) = \sin \frac{\pi x}{l}$.

Or

Solve the two-dimensional heat equation by the method of separation of variables.

- **5.** (a) Write the equation $3\frac{d^2x}{dt^2} + 6\frac{dx}{dt} x = t^2$ in normal form.
 - (b) Let $L \equiv D^2 + 2$, $f(t) = e^{2t}$, where $D \equiv \frac{d}{dt}$. Find Lf(t).

(Turn Over)

1

2

22P**/93**

(c) Transform the linear differential equation

$$\frac{d^2x}{dt^2} - 4\frac{dx}{dt} + 2x = t^2$$

into a system of first-order differential equation.

2

(d) Describe Euler's method.

4

Or

Find the characteristic roots of the equation associated in the solution of

$$\frac{dx}{dt} = 3x + y, \frac{dy}{dt} = 4x + 3y$$

(e) Solve:

$$\frac{dx}{dt} + \frac{dy}{dt} - x - 3y = e^{t}$$
$$\frac{dx}{dt} + \frac{dy}{dt} + x = e^{3t}$$

6

Or

Find $y(0\cdot 1)$, $y(0\cdot 2)$ in the solution of $\frac{dy}{dx} = x + y$, y(0) = 1, by using Runge-Kutta method.
