4 SEM TDC PHYH (CBCS) C 8

2022

(June/July)

PHYSICS

(Core)

Paper: C-8

(Mathematical Physics—III)

Full Marks: 53
Pass Marks: 21

Time: 3 hours

The figures in the margin indicate full marks for the questions

1. Choose the correct option:

 $1 \times 4 = 4$

- (a) If z_1 and z_2 are two complex numbers, then
 - (i) $|z_1+z_2| \ge |z_1|-|z_2|$
 - (ii) $|z_1 + z_2| \le |z_1| |z_2|$
 - (iii) $|z_1 + z_2| \le |z_1| |z_2| + |z_1 z_2|$
 - (iv) $|z_1 + z_2| \le |z_1| + |z_2| + |z_1 z_2|$

(b) The function
$$f(z) = \frac{1}{(z-2)^3}$$
 has a/a

an ____ at z = 2.

- (i) essential singularity
- (ii) pole
- (iii) branch point
- (iv) None of the above
- (c) The Laplace transform f(s) of F(t) = t is
 - (i) 1

(ii) s

(iii) s2

- (iv) $1/s^2$
- (d) If $g(\omega)$ is the Fourier transform of f(t), then the Fourier transform of f(at) is
 - (i) $\frac{1}{a}g\left(\frac{\omega}{a}\right)$
 - (ii) $\frac{1}{\omega}g\left(\frac{\omega}{a}\right)$
 - (iii) $\frac{1}{\omega}g\left(\frac{a}{\omega}\right)$
 - (iv) None of the above
- **2.** (a) Find the polar form of -5+5i.
- LAW.

2

(b) Find the residue of the function

$$f(z) = \frac{z}{(z-1)(z+1)^2}$$

- (c) Show how Cauchy's theorem can be used for a multiply connected region.
 - 2

(d) Show that the Fourier transform of the derivative of f(t) is $i\omega g(\omega)$, where $g(\omega)$ is the Fourier transform of f(t).

2

4

(e) Prove that if f(s) is the Laplace transform of F(t), then the Laplace transform of F(at) is

$$\frac{1}{a}f\left(\frac{s}{a}\right)$$
 2

3. (a) What are the different types of singularities of a complex function?

Locate and name the singularities of

$$f(z) = \frac{z^8 + z^4 + 2}{(z - 1)^3 (3z + 2)^2}$$
 2+3=5

(b) Prove Cauchy-Riemann equations in polar coordinates.

Or

If f(z) is an analytic function of z, then prove that

$$\left(\frac{\partial}{\partial x^2} + \frac{\partial}{\partial y^2}\right) |f(z)|^2 = 4|f'(z)|^2$$

(c) State the Cauchy's integral formula.

Evaluate

$$\oint_C \frac{\sin \pi z^2 + \cos \pi z^2}{(z-1)(z-2)} dz$$

where C is the circle |z|=1.

1+4=5

$$\int_{C} \frac{\sin^6 z}{\left(z - \frac{\pi}{6}\right)^3} dz$$

where C is the circle |z|=1.

4

3

(e) Express the following function in a Laurent's series:

 $f(z) = \frac{1}{(z+1)(z+3)}$

4. Find the Fourier transform of the following functions (any two): 3×2=6

- (i) $e^{-|t|}$
- (ii) $Ne^{-\alpha x^2}$ (N and α are constants)
- (iii) e^{-r^2/a^2} (a is a constant and $r = \sqrt{x^2 + y^2 + z^2}$)

- (i) $t^2 e^t \sin 4t$
- (ii) $e^{at}\cos\omega t$
- (iii) tn

3×2=6

- (a) Cauchy's theorem
- (b) Laplace transforms and its applications
- (c) Parseval's theorem
